

IRREDUCIBLE NONSURJECTIVE ENDOMORPHISMS OF F_n ARE HYPERBOLIC

JEAN PIERRE MUTANGUHA

ABSTRACT. We give a complete characterization of irreducible nonsurjective endomorphisms of F_n in terms of their topological representatives. Previously, Reynolds showed that any irreducible nonsurjective endomorphism can be represented by an irreducible immersion on a finite graph. We give a new proof of this and also show the converse holds when the immersion has connected Whitehead graphs. Consequently, irreducible nonsurjective endomorphisms are fully irreducible. The next result is a characterization of finitely generated subgroups of the free group that are invariant under an irreducible nonsurjective endomorphism. These two characterizations imply that the mapping torus of an irreducible nonsurjective endomorphism is word-hyperbolic.

1. INTRODUCTION

Bestvina-Handel defined irreducible automorphisms of free groups in [4] as the base case of their proof of the Scott conjecture. Irreducible automorphisms are the dynamical free group analogues of pseudo-Anosov diffeomorphisms of surfaces and, as such, the study of their dynamics has been an active area of research since their introduction. For instance, a leaf of the attracting lamination of a pseudo-Anosov diffeomorphism cannot have compact support in an infinite cover of the surface, and analogously, Bestvina-Feighn-Handel constructed a lamination for fully irreducible automorphisms and showed that a leaf of the lamination cannot be supported by a finitely generated infinite index subgroup [2, Proposition 2.4]. Another analogy: pseudo-Anosov mapping classes act with north-south dynamics on the Thurston compactification of Teichmüller space [22] and Levitt-Lustig proved that fully irreducible automorphisms act with north-south dynamics on the compactification of Culler-Vogtmann outer space [14].

We would like to extend these results to irreducible endomorphisms. We begin by defining the (semi)action of an injective endomorphism on outer space and the first result is that irreducible nonsurjective endomorphisms act with sink dynamics on outer space. There is a correspondence between having a fixed point of the action and being able to represent the endomorphism with a graph immersion. Under this correspondence, we prove:

Theorem 4.4. *If $\phi : F \rightarrow F$ is an irreducible nonsurjective endomorphism, then it can be represented by an irreducible immersion with connected Whitehead graphs. Moreover, the immersion is unique amongst irreducible immersions.*

All the terms in the theorem will be defined in Section 2. This is an unpublished result of Patrick Reynolds [18, Corollary 5.5] but the proof given here is original. We also note

that Reynolds' proof assumes the endomorphism is fully irreducible but, as we shall see shortly, this is equivalent to being irreducible. Not much is currently known about the action; for instance,

Problem 1.1. *Does the action of an irreducible nonsurjective endomorphism on outer space always have bounded image? See Example 3.3.*

Using the sink dynamics on outer space, we prove the converse of Theorem 4.4:

Theorem 4.5. *If $\phi : F \rightarrow F$ is represented by an irreducible immersion with connected Whitehead graphs, then ϕ is nonsurjective and fully irreducible.*

Dowdall-Kapovich-Leininger had previously shown the equivalence between the irreducible and fully irreducible properties for atoroidal automorphisms [10, Corollary B.4]. Theorem 4.4 and Theorem 4.5 give us this equivalence for nonsurjective endomorphisms.

Corollary 4.6. *Let $\phi : F \rightarrow F$ be an endomorphism. Then ϕ is nonsurjective and (fully) irreducible if and only if it is represented by an irreducible immersion with connected Whitehead graphs.*

The analogous criteria for fully irreducible automorphisms due to Catherine Pfaff [17, Theorem 4.1] and Ilya Kapovich [13, Theorem 1.2] require finding periodic Nielsen paths. So, in practice, the irreducibility of nonsurjective endomorphisms is easier to certify than that of automorphisms since it is straightforward to check that an immersion is irreducible and has connected Whitehead graphs. Furthermore, Theorem 4.5 can be combined with the action on the spine of outer space to get an algorithm that detects whether a nonsurjective endomorphism is irreducible. We sketch the algorithm at the end of Section 4.

The proof of the next proposition determines which finitely generated subgroups support a leaf of the lamination of an irreducible endomorphism; roughly speaking, these are the iterated images of the endomorphism, up to finite index.

Proposition 5.3. *Suppose $\phi : F \rightarrow F$ is injective and represented by an irreducible train track with connected Whitehead graphs. If $H \leq F$ is a finitely generated subgroup such that $\phi(H) \leq H$ and H contains a ϕ -expanding conjugacy class, then*

$$[\phi^k(F) : \phi^k(F) \cap H] < \infty \quad \text{for some } k \geq 0.$$

As a corollary of this technical result, we get a characterization for fully irreducible endomorphisms that applies to both automorphisms and nonsurjective endomorphisms.

Corollary 5.5. *Let $\phi : F \rightarrow F$ be an injective endomorphism. Then ϕ is fully irreducible if and only if $\phi : F \rightarrow F$ is primitively atoroidal, it is represented by an irreducible train track with connected Whitehead graphs, and its image $\phi(F)$ is not contained in a proper free factor.*

Another consequence of Proposition 5.3 is the hyperbolicity of the mapping torus of an irreducible immersion with connected Whitehead graphs – this was our original motivation.

Theorem 6.3. *If $\phi : F \rightarrow F$ is nonsurjective and irreducible, then $F *_{\phi}$ is word-hyperbolic.*

By our previous work [16], the proof boils down to showing that irreducible nonsurjective endomorphisms cannot have periodic laminations. As a quick application, we prove the hyperbolicity of the *Sapir group*:

Example 1.2. Let $F = F(a, b)$ be the free group on two generators and $\phi : F \rightarrow F$ be the nonsurjective endomorphism given by $\phi(a) = ab$ and $\phi(b) = ba$. The obvious map on the standard rose $f : R_2 \rightarrow R_2$ will be an immersion that induces ϕ on the fundamental groups. It is easy to verify that f is an irreducible immersion with a connected Whitehead graph. By Theorem 4.5, ϕ is fully irreducible, and by Theorem 6.3, the Sapir group $F*_\phi = \langle a, b, t \mid t^{-1}at = ab, t^{-1}bt = ba \rangle$ is word-hyperbolic.

In contrast, $\psi : F \rightarrow F$ given by $(a, b) \mapsto (aba, bab)$ is not word-hyperbolic: $\psi(ab) = (ab)^3$ and so $BS(1, 3) \cong \langle ab, t \rangle \leq F*_\psi$. As $H_1(F*_\phi) = \mathbb{Z}^2$ has rank > 1 , there are infinitely many isomorphisms $F*_\psi \cong F_n*_\psi$ [5]; all such endomorphisms $\psi_n : F_n \rightarrow F_n$ are nonsurjective, because $BS(1, 3) \leq F*_\psi$, and reducible by Theorem 6.3.

In future work, we use Proposition 5.3 to show that the irreducibility of an endomorphism is a group-invariant of its mapping torus, i.e., suppose $\phi : F_n \rightarrow F_n$ and $\psi : F_m \rightarrow F_m$ are endomorphisms (satisfying an additional necessary assumption for nonsurjective endomorphisms) and $F_n*_\phi \cong F_m*_\psi$, then ϕ is irreducible if and only if ψ is irreducible [15]. This answers Question 1.4 posed by Dowdall-Kapovich-Leininger in [11]; see also [10, Theorem C].

Outline: Section 2 contains the standard definitions and preliminary results that will be used throughout the paper. We define outer space and the action of injective endomorphisms on it in Section 3. Section 4 contains the proof of Theorem 4.4 and its converse. Theorem 4.5 then ends with an algorithm for detecting irreducible nonsurjective endomorphisms. In Section 5, we prove the technical result classifying subgroups that support a leaf of a lamination, Proposition 5.3. While the two previous sections are independent, we combine their main results in Section 6 to prove Theorem 6.3.

Acknowledgments: I would like to thank Ilya Kapovich for getting me started on the proof of Theorem 4.4. I also thank Derrick Wigglesworth and my advisor Matt Clay for the countless conversations on this material.

2. DEFINITIONS AND PRELIMINARY RESULTS

In this paper, F is a finitely generated free group on at least two generators.

Definition 2.1. An endomorphism $\phi : F \rightarrow F$ is **reducible** if there exists a free factorization $A_1 * \cdots * A_k * B$ of F , where B is possibly trivial if $k \geq 2$, and a sequence of elements, $(g_i)_{i=1}^k$, in F such that $\phi(A_i) \leq g_i A_{i+1} g_i^{-1}$ where the indices are considered mod k . An endomorphism ϕ is **irreducible** if it is not reducible and it is **fully irreducible** if all its iterates are irreducible; equivalently, ϕ is fully irreducible if there does not exist a proper free factor $A \leq F$, an element $g \in F$, and an integer $n \geq 1$ such that $\phi^n(A) \leq g A g^{-1}$.

An application of Stallings folds implies the endomorphisms studied in this paper will be injective even though it may not be explicitly stated again.

Lemma 2.2. *If $\phi : F \rightarrow F$ is irreducible, then it is injective.*

Proof. Stallings [20] showed that any endomorphism of F factors through folds whose nontrivial kernels corresponding to proper subgraphs, hence are normal closures of free factors. If ϕ is irreducible and not injective, then $\ker \phi$ contains an invariant free factor; irreducibility implies the only invariant free factor is the group itself, but this implies ϕ is trivial, in which case, all proper free factors are also invariant – a contradiction. \square

Definition 2.3. Fix an isomorphism $F \cong \pi_1(\Gamma)$ for some connected finite core graph Γ , i.e., a finite 1-dimensional CW-complex with no univalent vertices. For any subgroup $H \leq F$, the **Stallings subgroup graph** $S(H)$ is the core of the cover of Γ corresponding to H , i.e., it is the smallest subgraph of the cover containing all immersed loops of the cover. When H is nontrivial and finitely generated, $S(H)$ is a finite core graph with rank ≥ 1 . A Stallings subgroup graph comes with an immersion $v : S(H) \rightarrow \Gamma$, which is a restriction of the covering map. We shall also assume that $S(H)$ has been subdivided so that v is **simplicial**: v maps edges to edges. The immersion $S(H) \rightarrow \Gamma$ uniquely determines $[H]$, the conjugacy class of H : if there is a subgroup $H' \leq F$ with immersion $v' : S(H') \rightarrow \Gamma$ and homeomorphism $h : S(H) \rightarrow S(H')$ such that $v = v' \circ h$, then $[H] = [H']$.

The last statement makes Stallings subgroup graphs especially useful for studying a nonsurjective injective endomorphism $\phi : F \rightarrow F$. For $i \geq 1$, let $S_i = S(\phi^i(F))$. When ϕ is an automorphism, the maps $v_i : S_i \rightarrow \Gamma$ are all graph isomorphisms. Conversely, suppose there was a homeomorphism $h : S_i \rightarrow S_j$ for some $i < j$ such that $v_i = v_j \circ h$. Then, by the last statement of the definition, $[\phi^i(F)] = [\phi^j(F)]$. This means there is some element $g \in F$ such that $g\phi^i(F)g^{-1} = \phi^j(F) \leq \phi^i(F)$. But in free groups (subgroup separable), a finitely generated subgroup cannot be conjugate to a proper subgroup. Hence $\phi^j(F) = \phi^i(F)$. But $\phi^{j-i}|_{\phi^i(F)}$ is conjugate to ϕ^{j-i} as ϕ is injective. In particular, ϕ^{j-i} is an automorphism and so is ϕ . Therefore, when ϕ is nonsurjective, the maps v_i are all distinct and the number of edges in S_i grows as $i \rightarrow \infty$. We shall use this fact in Section 4 to show that, for an irreducible nonsurjective endomorphism, the sequence of maps $S_i \rightarrow S_i$ that induce ϕ on $\pi_1(S_i) \cong F$ converges to an immersion on Γ that induces ϕ on $\pi_1(\Gamma) \cong F$.

Definition 2.4. Let Γ be a finite graph. A map $f : \Gamma \rightarrow \Gamma$ is a **train track** if it maps vertices to vertices and all of its iterates are immersions when restricted to the interior of any edge. A train track is **irreducible** if for any pair of edges e_i, e_j in Γ , e_i is in the image of some f -iterate of e_j . Given a train track $f : \Gamma \rightarrow \Gamma$, we fix an ordering of the edges of Γ and construct the **transition matrix** $A(f)$ as follows: it is a nonnegative square matrix whose size is given by the number of edges in Γ ; the (i, j) -th entry of $A(f)$ is the number of times the edge e_i appears in the image of e_j . An irreducible train track is **expanding** if it is not a homeomorphism; equivalently, the transition matrix is *irreducible* and has a real eigenvalue $\lambda_f > 1$ – this is the *Perron-Frobenius eigenvalue*.

Let $f : \Gamma \rightarrow \Gamma$ be a train track map and v be a vertex of Γ . The **Whitehead graph** at v is a simple graph whose vertices are the half-edges of Γ attached to v , denoted by $T_v(\Gamma)$. A

pair of elements of $T_v(\Gamma)$ is called a **turn** at v and it is **nondegenerate** if the pair consists of distinct elements. A nondegenerate turn is **f -legal** if it remains nondegenerate under iteration of f . A turn at v is an edge of the Whitehead graph if it appears in the image of an f -iterate of some edge of Γ . Note that the edges in the Whitehead graphs are f -legal.

An irreducible train track $f : \Gamma \rightarrow \Gamma$ is **weakly clean** if the Whitehead graph at each vertex is connected. A weakly clean map is **clean** if there is an iterate such that every edge surjects onto the whole graph; equivalently, the transition matrix is *primitive*. It follows from the definition that if a map f is clean then so are all its iterates f^i ($i \geq 1$).

The following proposition due to Dowdall-Kapovich-Leininger [10, Proposition B.2] allows us to use clean and weakly clean interchangeably. We give a proof that does not assume homotopy-equivalence.

Proposition 2.5. *Suppose Γ is a connected graph and $f : \Gamma \rightarrow \Gamma$ is a train track with an irreducible but not primitive transition matrix. Then f has a vertex with a disconnected Whitehead graph. In particular, if f is weakly clean, then f is clean.*

Proof. As $A(f)$ is irreducible but not primitive, it is permutation-similar to a transitive block permutation matrix [1, Theorem 1.8.3]. This block permutation form gives a partition of $E(\Gamma)$ into $d \geq 2$ proper subgraphs $\Gamma_0, \dots, \Gamma_{d-1}$ such that $f(\Gamma_i) \subset \Gamma_{i+1}$ where indices are considered mod d . Thus, any vertex adjacent to two or more of these subgraphs has a disconnected Whitehead graph. Such a vertex exists since Γ is connected. \square

Definition 2.6. Two homomorphisms $\phi_1, \phi_2 : A \rightarrow B$ are *equivalent* if there is an inner automorphism of B , i_g , such that $\phi_1 = i_g \circ \phi_2$. An *outermorphism* is an equivalence class in $\text{Hom}(A, B)$, denoted by $[\phi]$, and $\text{Out}(F)$ is the group of outer automorphisms of F . A graph map $f : \Gamma \rightarrow \Gamma$ is a **(topological) representative** for an injective (outer) endomorphism $\phi : F \rightarrow F$ if: Γ is a connected core graph; the map f maps vertices to vertices and is locally injective in interior of edges and at bivalent vertices; and there is an isomorphism $\alpha : F \rightarrow \pi_1(\Gamma)$, known as a **marking**, such that $[f_*\alpha] = [\alpha\phi]$.

Bestvina-Handel defined train tracks in [4] and one of the main results was the algorithmic construction of train track representatives for irreducible endomorphisms. We note that their result was stated for irreducible automorphisms but the proof itself never used nor needed the fact that the endomorphisms were surjective. See also Dicks-Ventura [9].

Theorem 2.7 ([4, Theorem 1.7]). *If $\phi : F \rightarrow F$ is an irreducible endomorphism, then ϕ can be represented by an irreducible train track map. The irreducible train track is expanding if and only if ϕ has infinite-order.*

The following argument is now a standard technique in the field and has its roots in Bestvina-Handel's paper [4, Proposition 4.5].

Corollary 2.8. *If $\phi : F \rightarrow F$ is irreducible and has infinite-order, then ϕ has a clean representative. Moreover, any irreducible train track representative of ϕ will be clean.*

Sketch proof. By Theorem 2.7, the endomorphism ϕ has an expanding irreducible train track representative. If the Whitehead graph of some vertex were disconnected, then

blowing-up the vertex and the appropriate preimages to separate the components of the Whitehead graph would give a reduction of ϕ . So all Whitehead graphs are connected and the representative is (weakly) clean. For more details, see [12, Proposition 4.1] \square

Notation. We denote by $[p]$ (resp. $[\rho]$) the immersed path (resp. loop) homotopic rel. endpoints to a path p (resp. freely homotopic to a loop ρ).

For any irreducible train track map $f : \Gamma \rightarrow \Gamma$, there is the associated Perron-Frobenius eigenvalue of the transition matrix, $\lambda_f \geq 1$, and a metric on the graph Γ given by the Perron-Frobenius left eigenvector. With this metric, f is homotopic rel. vertices to a (unique) map whose restrictions to edges are local λ_f -homotheties. For the rest of the paper, we shall assume an irreducible train track map has the latter linear property. The following lemma, known as the **Bounded Cancellation Lemma**, combined with the train track linear structure allow us to carefully study the dynamics of f .

Lemma 2.9 ([9, Lemma II.2.4]). *Suppose $f : \Gamma \rightarrow \Gamma$ is a topological representative for an injective endomorphism $\phi : F \rightarrow F$. Then there exists an constant $C = C(f)$ such that for any immersed path that can be written as a concatenation $a \cdot b$ in Γ , there exists path decompositions $[f(a)] = x \cdot u$, $[f(b)] = \bar{u} \cdot y$, $[f(a \cdot b)] = x \cdot y$, where the length of $u < C$.*

Suppose $f : \Gamma \rightarrow \Gamma$ is an expanding irreducible train track and let $C = C(f)$ be the constant given by the bounded cancellation lemma. If $a \cdot b \cdot c$ is an immersed path and b is f -legal, then $f(a) = x \cdot u$, $f(b) = \bar{u} \cdot y \cdot v$, $f(c) = \bar{v} \cdot z$ and $[f(a \cdot b \cdot c)] = x \cdot y \cdot z$ where the lengths of u and v are less than C and y is f -legal. With the assumed linear structure on train tracks, we have $l(y) = \lambda_f \cdot l(b) - l(u) - l(v) > \lambda_f \cdot l(b) - 2C$. A quick calculation, shows that if $l(b) > \frac{2C}{\lambda_f - 1}$, then there is a nontrivial subpath s of b such that $f^k(s)$ is a subpath of $[f^k(a \cdot b \cdot c)]$ for any $k \geq 1$ and immersed path $a \cdot b \cdot c$. The number $\frac{2C}{\lambda_f - 1}$ is known as the **critical constant**.

3. THE ACTION OF INJECTIVE ENDOMORPHISMS ON OUTER SPACE

Culler-Vogtmann introduced outer space in [8] as a topological space with a nice $\text{Out}(F)$ -action. We will give two equivalent definitions and then describe the $\text{Out}(F)$ -action and, more generally, the semi-action of an injective endomorphism with respect to the two descriptions of outer space. See Karen Vogtmann's survey paper [23] for details.

Definition 3.1. In general, **trees** refers to *real trees* (0-hyperbolic geodesic spaces) and a tree is **simplicial** if it is homeomorphic to a locally finite CW-complex. The **Culler-Vogtmann outer space** of F , denoted by $CV(F)$, is the set of connected simplicial trees, T , with a free minimal (left) action of F , $\alpha : F \rightarrow \text{Isom}(T)$, up to the equivalence: $(T_1, \alpha_1) \sim (T_2, \alpha_2)$ if there is an F -equivariant surjective homothety $f : T_1 \rightarrow T_2$, i.e., $f \circ \alpha_1(g) = \alpha_2(g) \circ f$ for all $g \in F$. Alternatively, $CV(F)$ is the set of connected finite metric core graphs, Γ , with a marking $\alpha : F \rightarrow \pi_1(\Gamma)$, up to the equivalence: $(\Gamma_1, \alpha_1) \sim (\Gamma_2, \alpha_2)$ if there is a homothety $f : \Gamma_1 \rightarrow \Gamma_2$ such that $[f_* \alpha_1] = [\alpha_2]$. Identify $\pi_1(\Gamma)$ with the group of deck transformations of $\tilde{\Gamma}$ to get the correspondence between the two descriptions.

We will now equip $CV(F)$ with a topology: for an equivalence class $[\Gamma, \alpha] \in CV(F)$, choose a representative such that Γ has no bivalent vertices and $\text{vol}(\Gamma) = 1$; the volume of a metric graph $\text{vol}(\Gamma)$ is the sum of the lengths of all edges in the graph. Let n be the number of edges in Γ and identify $\sigma(\Gamma, \alpha)$ with the $(n - 1)$ -simplex one gets by varying the lengths of the edges of Γ to get homeomorphic Γ' while still maintaining the equality $\text{vol}(\Gamma') = 1$. In the tree description of $CV(F)$, this variation corresponds to equivariantly varying the metric on $(\tilde{\Gamma}, \alpha)$ to get $(\tilde{\Gamma}', \alpha')$.

This gives us a decomposition of $CV(F)$ into a disjoint union of open simplices $\sigma(\Gamma, \alpha)$. Attaching maps for these simplices are given by decreasing the volume of some forest of Γ to 0. This decomposition of $CV(F)$ and description of attaching maps makes $CV(F)$ a locally finite *open simplicial complex*, i.e., a simplicial complex with some missing faces corresponding to collapsing noncontractible subgraphs. The set of open simplices of $CV(F)$, denoted by $SCV(F)$, can be made into a locally finite simplicial complex known as the *spine of outer space* but, for the most part, we will treat it as a set with no added structure.

For the rest of the paper, we will assume $CV(F)$ has this topology. It is a theorem of Culler-Vogtmann that $CV(F)$ is contractible [8, 24]. There is a natural right action of $\text{Out}(F)$ on $CV(F)$ given by $[T, \alpha] \cdot [\phi] = [T, \alpha\phi]$ for $[T, \alpha] \in CV(F)$ and $[\phi] \in \text{Out}(F)$. Furthermore, this action induces an action on $SCV(F)$: $\sigma(T, \alpha) \cdot [\phi] = \sigma(T, \alpha\phi)$. For an injective (outer) endomorphism $\phi : F \rightarrow F$, there is a right (semi)action given by $[T, \alpha] \cdot [\phi] = [T', \alpha\phi]$ where T' is the minimal tree of the *twisted* action $\alpha\phi : F \rightarrow \text{Isom}(T)$, i.e., T' is the minimal tree for $\phi(F)$. This induces an action on $SCV(F)$ by forgetting the metrics: $\sigma(T, \alpha) \cdot [\phi] = \sigma(T', \alpha\phi)$. For an F -tree (T, α) and a marked metric graph (Γ, α) , we shall abuse notation and write only T and Γ when the actions and markings are clear. The goal of the next section is to describe dynamics of the action of irreducible nonsurjective endomorphisms on $CV(F)$. We will show that the action has a unique attracting point and no other fixed points.

We now give the second description of the action of an injective endomorphism $\phi : F \rightarrow F$ on $CV(F)$ and $SCV(F)$. Let Γ be a marked metric graph, i.e., $[\Gamma, \alpha] \in CV(F)$. Fix any topological representative $f : \Gamma \rightarrow \Gamma$ for ϕ . For any $i \geq 1$, the map $f^i : \Gamma \rightarrow \Gamma$ factors as $f^i = v_i h_i$ where $h_i : \Gamma \rightarrow S'_i$ is a composition of folds and $v_i : S'_i \rightarrow \Gamma$ is an immersion. The graph S'_i is subdivided so that the immersion is simplicial. If we let $S_i \subset S'_i$ be the core subgraph, then this is the Stallings subgroup graph for $\phi^i(F)$ and the graphs fits in the following commutative diagram:

$$\begin{array}{ccccccc}
 & & S_1 & & S_i & & S_{i+1} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \Gamma & \longrightarrow & S'_1 & \longrightarrow & \cdots & \longrightarrow & S'_i & \longrightarrow & S'_{i+1} & \longrightarrow & \cdots \\
 \parallel & & \downarrow v_1 & & \downarrow v_i & & \downarrow v_{i+1} & & & & \\
 \Gamma & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \cdots & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \cdots
 \end{array}$$

Since ϕ is injective, the composition of folds $h_i : \Gamma \rightarrow S'_i$ is a homotopy equivalence which induces a marking $h_{i*}\alpha : F \rightarrow \pi_1(S'_i) = \pi_1(S_i)$. Pullback the metric on Γ via the immersion $v_i : S_i \rightarrow \Gamma$ to get a metric on S_i . By construction, $\tilde{v}_i(\tilde{S}_i) \subset \tilde{\Gamma}$ is the minimal tree for $\phi^i(F)$ and $[\tilde{\Gamma}] \cdot [\phi]^i = [\tilde{S}_i]$ as \tilde{v}_i is an isometric embedding. Therefore, in terms of marked metric graphs, the action is given by $[\Gamma, \alpha] \cdot [\phi]^i = [S_i, h_{i*}\alpha]$. As the immersion v_i and folds h_i do not depend on the metric on Γ , we see that the action of ϕ on $CV(F)$ is piecewise-linear with respect to the open simplicial structure and it induces the action on $SCV(F)$ by forgetting the metrics.

The following lemma tells us precisely when ϕ fixes an element of $SCV(F)$ or $CV(F)$.

Lemma 3.2. *Let $\phi : F \rightarrow F$ be an injective endomorphism. Then:*

- (1) $\sigma(\Gamma) \in SCV(F)$ is fixed by ϕ if and only if ϕ is represented by an immersion on Γ .
- (2) $[\Gamma] \in CV(F)$ is fixed by ϕ if and only if ϕ is represented by a local homothety on Γ .

Proof. Suppose $\sigma(\Gamma, \alpha) \cdot [\phi] = \sigma(\Gamma, \alpha)$, i.e., $\sigma(S_1, h_{i*}\alpha) = \sigma(\Gamma, \alpha)$. Then the composition of folds and core retraction $h_1 : \Gamma \rightarrow S_1$ is homotopic to a homeomorphism. As $f = v_1 h_1$ and v_1 is an immersion, we get that f is homotopic to an immersion. Suppose $[\Gamma, \alpha] \cdot [\phi] = [\Gamma, \alpha]$. By the previous sentence, we may assume f is an immersion, $S_1 = S'_1$, and $h_1 : \Gamma \rightarrow S_1$ is a homeomorphism. Finally, for the point $[\Gamma, \alpha]$ to be fixed, the immersion $v_1 : S_1 \rightarrow \Gamma$ and homeomorphism $h_1 : \Gamma \rightarrow S_1$ must induce the same projective metric on S_1 . Thus f is (homotopic to) a local homothety. \square

We end the section with a description of the action of the endomorphism in Example 1.2.

Example 3.3. Recall $F = F(a, b)$ and $\phi : F \rightarrow F$ is given by $(a, b) \mapsto (ab, ba)$. In rank 2, the spine of outer space has the structure of a regular trivalent tree with a spike at the midpoint of edges. Set $F = \pi_1(R_2)$, then the standard rose is the marked graph $R_* = (R_2, id_F)$ and all other roses are given by (R_2, φ) for some $[\varphi] \in \text{Out}(F)$. Let (B, β) be the marked barbell graph attached to R_* and let (T, θ) be the theta graph between R_* , (R_2, φ_a) , and (R_2, φ_b) where $\varphi_a : (a, ab)$ and $\varphi_b : (ba, b)$ are the generators of $\text{Out}(F)$.

To compute $\sigma(R_2, \varphi) \cdot [\phi]$, first represent $\varphi\phi$ on R_* then fold. The composition of folds gives the resulting marked graph. For instance, $\sigma(R_2, \varphi_a) \cdot [\phi] = \sigma(T, \theta) = \sigma(R_2, \varphi_b) \cdot [\phi]$, $\sigma(R_2, \varphi_a^{-1}) \cdot [\phi] = \sigma(B, \beta)$, and $\sigma(R_2, \varphi_a^{-1}) \cdot [\phi]^2 = \sigma(T, \theta\varphi_a^{-1})$. Figure 1 below illustrates one of these computations. Along these lines, we can show that


$$S = \{\sigma(R_*), \sigma(T, \theta), \sigma(T, \theta\varphi_a^{-1}), \sigma(B, \beta)\}$$

is the set of ϕ -periodic elements in $SCV(F)$; the first two are ϕ -fixed while the latter two have ϕ -period 2. By inducting on the roses of the spine, we can verify $SCV(F) \cdot [\phi] = S$.

4. IRREDUCIBLE NONSURJECTIVE ENDOMORPHISMS ARE IMMERSIONS

Given an expanding irreducible train track map $f : \Gamma \rightarrow \Gamma$ representing an injective endomorphism $\phi : F \rightarrow F$, then ϕ has a right action on the F -tree $\tilde{\Gamma}$ by taking minimal trees of the twisted actions: $(\tilde{\Gamma}, \alpha) \cdot \phi^n = (\tilde{\Gamma}', \alpha\phi^n)$. We define the limit tree

$$T_f = \lim_{n \rightarrow \infty} \lambda_f^{-n} \tilde{\Gamma} \phi^n$$


 FIGURE 1. Computation showing $\sigma(R_2, \varphi_a^{-1}) \cdot [\phi] = \sigma(B, \beta)$.

where the limit is taken in the space of length functions on F .

Bestvina-Feighn-Handel [2] show that $[T] \cdot [\phi]^n \rightarrow [T_f]$ for any $[T] \in CV(F)$; here the limit is taken in the space of projective length functions. The class $[T_f]$ need not be in $CV(F)$ and, in fact, it is not when ϕ is an automorphism. We will use $[T_\phi] = [T_f]$ and $\lambda_\phi = \lambda_f$ to emphasize the independence from the choice of train track representative f . The F -action on T_ϕ is free if and only if ϕ is atoroidal, i.e., it has no periodic conjugacy class. When ϕ is represented by an irreducible immersion $f : \Gamma \rightarrow \Gamma$ where Γ has the train track linear structure, then f is a local λ_ϕ -homothety, $T_\phi = \hat{\Gamma}$, and $[T_\phi] \in CV(F)$. In a certain sense, the immersion is unique since $[\Gamma]$ is the unique ϕ -fixed point in $CV(F)$.

In the general setting considered in the previous section, not much can be said about the graphs S_i and immersions $v_i : S'_i \rightarrow \Gamma$. However, if $f : \Gamma \rightarrow \Gamma$ is assumed to be a clean representative, we gain more control; for example, we will show in this case that S'_i is a core graph, hence, $S_i = S'_i$. To do this, we need to introduce a new structure:

Definition 4.1. Let $f : \Gamma \rightarrow \Gamma$ be an irreducible train track map and, for some $i \geq 1$, $f^i = hg$ where $g : \Gamma \rightarrow X$ and $h : X \rightarrow \Gamma$ are surjective graph maps and X is a finite core graph. A turn at a vertex v of X is **legal** if its image under h is an f -legal turn. The **relative Whitehead graph** of (X, g, h) at v is a simple graph whose vertex set is $T_v(X)$ and a turn at v is an edge of the relative Whitehead graph if it is in the edge-path $g(f^j(e))$ for some $j \geq 0$ and edge e in Γ . Note that edges of the relative Whitehead graphs are legal.

The next lemma tells us how subdivision and folding affect the relative Whitehead graphs.

Lemma 4.2. Suppose $f^i : \Gamma \xrightarrow{g} X \xrightarrow{h} \Gamma$ has connected relative Whitehead graphs and $h = h_2 h_1$ where $h_1 : X \rightarrow Y$ is a subdivision or a single fold. Then Y is a core graph and the relative Whitehead graphs of $(Y, h_1 g, h_2)$ are connected.

Proof. Suppose h_1 is a subdivision. If v is a vertex of Y that was added by the subdivision, then $T_v(Y)$ has two elements, $v \in \text{Int}(e)$ for some edge e' of X , and $v \in h_1(e')$. In particular, the two elements of $T_v(Y)$ are joined by an edge since g is surjective.

If v is a vertex of both Y and X , then $T_v(Y) = T_v(X)$ and subdivision makes no changes to the relative Whitehead graph at v . So the relative Whitehead graph of $(Y, h_1 g, h_2)$ is connected since we assumed (X, g, h) has connected relative Whitehead graphs.

Suppose h_1 is a single fold. The only way to produce a univalent vertex in Y with a fold is by folding a bivalent vertex of X . But bivalent vertices only have one turn which is legal since the relative Whitehead graphs of (X, g, h) are connected.

At the origin vertex, the fold identifies two distinct vertices of the relative Whitehead graphs and this preserves connectedness. At the two terminal vertices, the fold identifies two distinct vertices, one from each relative Whitehead graph, and this gluing of two connected graphs produces a connected graph. If a vertex in X is not part of the fold, then its image in Y will have the same connected relative Whitehead graph. \square

Recall that to construct S'_i , we subdivide and fold the map $f^i : \Gamma \rightarrow \Gamma$ to get the immersion $v_i : S'_i \rightarrow \Gamma$. Suppose f was a clean map. For the base case, the relative Whitehead graphs of (Γ, id_Γ, f^i) are connected as they are isomorphic to the Whitehead graphs of f^i . By Lemma 4.2 and induction on folds, S'_i is a core graph and the relative Whitehead graphs of (S_i, h_i, v_i) are connected. So the immersion v_i maps edge-paths connecting branch points (*natural edges*) of S_i to f -legal edge-paths in Γ . The commutative diagram in the previous section becomes:

$$\begin{array}{ccccccccccc} \Gamma & \longrightarrow & S_1 & \longrightarrow & S_2 & \longrightarrow & \cdots & \longrightarrow & S_i & \longrightarrow & S_{i+1} & \longrightarrow & \cdots \\ \parallel & & \downarrow v_1 & & \downarrow v_2 & & & & \downarrow v_i & & \downarrow v_{i+1} & & \\ \Gamma & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \cdots & \xrightarrow{f} & \Gamma & \xrightarrow{f} & \Gamma & \longrightarrow & \cdots \end{array}$$

Aside from the construction of this diagram and the next lemma, this section and the next are independent of each other. This lemma will only be used in the proof of Proposition 5.3, the main result of the next section.

Lemma 4.3. *Let f, v_i, h_i be as in the previous paragraph. The map $\hat{f}^i : S_i \rightarrow S_i$ given by $\hat{f}^i = h_i v_i$ is a clean representative for ϕ^i .*

Proof. For all $n \geq 1$, $(\hat{f}^i)^{n+1} = (h_i v_i)^{n+1} = h_i (v_i h_i)^n v_i = h_i f^{in} v_i$. Since f is a train track, v_i is a simplicial immersion, and h_i folds illegal turns only, it follows that \hat{f}^i is a train track too. Since f is clean, there is an $n \geq 1$ such that, for any edge e in Γ , $f^{in}(e)$ is surjective and contains all the edges of the Whitehead graphs of f^i . As h_i is surjective, $(\hat{f}^i)^{n+1}$ is surjective when restricted to any edge and \hat{f}^i has a primitive transition matrix. Since folding only identifies vertices of relative Whitehead graphs, $h_i f^{in}(e)$ contains all the edges of the relative Whitehead graphs of (S_i, h_i, v_i) . As v_i maps edges to edges, we have that the Whitehead graphs of \hat{f}^i contains the relative Whitehead graphs of (S_i, h_i, v_i) as subgraphs. Since v_i is surjective, we have the reverse containment too. So the Whitehead graphs of \hat{f}^i are connected and the map is clean. \square

With this set-up, we are now ready to present a new proof of an unpublished theorem due to Patrick Reynolds [18, Corollary 5.5]. The first half of this proof is based on an argument due to Ilya Kapovich showing that irreducible nonsurjective endomorphisms are expansive; c.f. [18, Proposition 3.11].

Theorem 4.4. *If $\phi : F \rightarrow F$ is an irreducible nonsurjective endomorphism, then it can be represented by a clean immersion, unique amongst irreducible immersions.*

Proof. Let $f : \Gamma \rightarrow \Gamma$ be a clean representative for $\phi : F \rightarrow F$ given by Corollary 2.8. By the discussion following Lemma 4.2, we now construct core graphs S_i , folds $h_i : \Gamma \rightarrow S_i$, and immersions $v_i : S_i \rightarrow \Gamma$ such that $f^i = v_i h_i$ and the natural edges of S_i are legal, i.e., they consist of legal turns. Nonsurjectivity implies that the number of edges in S_i grows as $i \rightarrow \infty$. So for some $i \gg 0$, S_i has a natural edge longer than the critical constant. By the bounded cancellation lemma, there is a nontrivial subpath s of the long natural edge such that, for all $j > i$, the folds in $S_i \rightarrow S_j$ are supported in the complement of s .

Suppose $i \gg 0$ is an index such that S_i has the maximal number of natural edges longer than $\frac{2C \cdot L \cdot \lambda_\phi}{\lambda_\phi - 1}$, where $L = L(\text{rank}(F))$ is the maximum number of natural edges of an embedded loop of a marked graph. Denote by K_i the proper subgraph consisting of the remaining *short* natural edges of S_i with length at most the critical constant. Let $\hat{f} : S_i \rightarrow S_i$ be a map representing ϕ , i.e., $[\hat{f}_* h_{i*}] = [h_{i*} f_*]$. Suppose K_i has an embedded loop ρ , then by construction $f v_i(\rho)$ decomposes into at most L legal paths, each of length at most $\frac{2C \lambda_\phi}{\lambda_\phi - 1}$. So $[f v_i(\rho)]$ lifts to the loop $[\hat{f}(\rho)]$ in K_i . Thus if K_i is not a forest, then K_i is an \hat{f}_i -invariant proper subgraph up to homotopy. Since ϕ is irreducible, K_i must be a forest. Therefore, the loops in S_i are growing exponentially, ϕ is atoroidal, and the action of F on T_ϕ is free.

We showed in the previous paragraph that folds in the map $S_i \rightarrow S_j$ ($j > i$) are supported in some forest $K \subset S_i$. Since there are finitely many combinatorially distinct ways to fold a forest, there is a $j > i$ and $k \geq 1$ such that the composition of folds $S_j \rightarrow S_{j+k}$ is homotopic to a homeomorphism. This means $[\tilde{S}_j], [\tilde{S}_{j+k}]$ lie in the simplex $\sigma(\tilde{S}_j) \subset CV(F)$. In particular, $\sigma(\tilde{S}_j) \cdot [\phi]^k = \sigma(\tilde{S}_j)$ and the sequence $[\tilde{S}_{j+kn}]$ ($n \geq 0$) lies in $\sigma(\tilde{S}_j)$.

So the projective limit tree for this sequence,

$$\lim_{n \rightarrow \infty} [\tilde{S}_{j+kn}] = \lim_{n \rightarrow \infty} [\tilde{S}_j] \cdot [\phi]^{kn} = \lim_{n \rightarrow \infty} [\tilde{S}_j] \cdot [\phi]^n = [T_\phi],$$

is a simplicial tree, i.e., the F -quotient of T_ϕ , call it Γ_0 , is a graph obtained by collapsing a (possibly empty) subgraph of S_j . As T_ϕ is free and simplicial, $[T_\phi]$ is a point in $CV(F)$. By definition and Lemma 3.2, $[T_\phi]$ is fixed by ϕ and ϕ is represented by a local λ_ϕ -homothety $f_0 : \Gamma_0 \rightarrow \Gamma_0$. Furthermore, the local homothety f_0 is expanding since $\lambda_\phi > 1$; so f_0 has no invariant forests and the irreducibility of ϕ implies f_0 is clean by Corollary 2.8. Uniqueness follows from Lemma 3.2 and uniqueness of the (attracting) fixed point $[T_\phi]$. \square

This theorem tells us that the action of an irreducible nonsurjective endomorphism on $CV(F)$ has a unique fixed point that is also a global attracting point. Reynolds studied this action further; for instance, the action converges to the fixed point uniformly on compact sets and, for *admissible* endomorphisms, the action can be extended to the compactification of outer space $\overline{CV(F)}$ where the convergence is uniform.

We now use the action on outer space again to prove the converse to Theorem 4.4, which can also be thought of as a criterion for fully irreducible nonsurjective endomorphisms.

Theorem 4.5. *If $\phi : F \rightarrow F$ is represented by a clean immersion, then ϕ is nonsurjective and fully irreducible.*

Proof. Since ϕ is represented by a clean immersion $f : \Gamma \rightarrow \Gamma$, it is nonsurjective. Suppose $\phi^n(A) \leq gAg^{-1}$ where $A \leq F$ is a proper free factor, $g \in F$, and $n \geq 1$. Let $S(A)$ be the Stallings subgroup graph corresponding to A with respect to Γ . Set $\psi = i_g \circ \phi^n$ so that $\psi(A) \leq A$ and the immersion f^n lifts to an immersion $g : S(A) \rightarrow S(A)$ representing $\psi|_A$. Complete $\Delta_0 = S(A)$ to a graph Γ' with a marking $\pi_1(\Gamma') \cong F$ and extend g to the rest of Γ' such that $g : \Gamma' \rightarrow \Gamma'$ is a topological representative for ψ and Δ_0 is a noncontractible g -invariant proper subgraph corresponding to A .

Recall from Section 3, the Stallings subgroup graph $S_i = S(\psi^i(F))$ with respect to Γ' along with the marking given by folding $g^i : \Gamma' \rightarrow \Gamma'$ determine the i -th iterate of $[\Gamma'] \in CV(F)$ under the ψ -action, i.e., $[\Gamma'] \cdot [\psi]^i = [S_i]$. Since $[S_i] \rightarrow [\Gamma]$, as discussed at the start of this section, and the spine of outer space is locally finite, we have that the sequence $\sigma(S_i)$ ($i \geq 1$) in $SCV(F)$ is eventually periodic. So for some $i, j \geq 1$, we have $\sigma(S_i) = \sigma(S_{i+j})$ and, by Lemma 3.2, ψ^j can be represented by an immersion on S_i .

Let $h_i : \Gamma' \rightarrow S'_i$ be the composition of folds given by the construction of S_i . Since $g|_{\Delta_0}$ is an immersion, the corresponding restriction $h_i|_{\Delta_0}$ is either a homeomorphism or an identification of vertices of Δ_0 . Set $\Delta_i = h_i(\Delta_0)$ and let $g^j_i : S_i \rightarrow S_i$ be the induced map representing ψ^j . Then Δ_i is a noncontractible g_i -invariant subgraph and $g_i|_{\Delta_i}$ is an immersion. By the previous paragraph, g^j_i is homotopic to an immersion $\gamma : S_i \rightarrow S_i$. The homotopy will preserve the invariance of Δ_i so that Δ_i is a γ -invariant subgraph. As $\sigma(S_i)$ is fixed by ψ^j , the sequence $[S_{i+jm}]$ is constructed by pulling back the metric of S_i via γ^m .

Iteratively pulling back the metric via γ has the effect of collapsing γ -invariant forests and normalizing the metric so the induced map is a local homothety. By uniqueness of the limit $[S_{i+jm}] \rightarrow [\Gamma]$, the induced map must be f^{nj} and Δ_∞ , the image of Δ_i under the collapse map, is an f^{nj} -invariant subgraph. But f^{nj} is a clean map, so $\Delta_\infty = \Gamma$.

Let $h_\infty : \Delta_0 \rightarrow \Delta_\infty = \Gamma$ be the induced map. By construction, $f^{nj} \circ h_\infty = h_\infty \circ g^j|_{\Delta_0}$ and, as h_∞ is an identification of some vertices and possibly a collapse of a forest, the Whitehead graphs of f^{nj} are determined by where g^j maps the edges of Δ_0 . So f^{nj} will have disconnected Whitehead graphs at the identified vertices – a contradiction. \square

A result due to Dowdall-Kapovich-Leininger is that atoroidal irreducible automorphisms are fully irreducible [10, Corollary B.4]. Combining Theorem 4.4 and Theorem 4.5, we get that this holds for irreducible nonsurjective endomorphisms.

Corollary 4.6. *Let $\phi : F \rightarrow F$ be an endomorphism. Then ϕ is nonsurjective and (fully) irreducible if and only if it is represented by a clean immersion.*

In the next section, we shall give a uniform characterization for fully irreducible endomorphism that holds for both surjective and nonsurjective endomorphisms. We now sketch an algorithm for checking whether a nonsurjective endomorphism is irreducible.

Let $\phi : F \rightarrow F$ be a given nonsurjective endomorphism:

- (1) Check if the Stallings subgroup graph $S(\phi(F))$ with respect to the standard rose has the same rank as F . If not, then ϕ is not injective and we are done by Lemma 2.2.
- (2) Run Bestvina-Handel's algorithm to construct an irreducible train track representative [4, Theorem 1.7]. This algorithm either succeeds or produces a reduction for ϕ .

In the latter case, we are done and ϕ is reducible. Otherwise, assume $f : \Gamma \rightarrow \Gamma$ is an irreducible train track representative for ϕ .

- (3) By Corollary 2.8, either the Whitehead graphs of f are connected or there is a reduction for ϕ . In the latter case, we are done. Otherwise, assume f is clean.
- (4) Construct graphs $S_i = S(\phi^i(F))$ with respect to Γ by folding iterates of f . Let $f_i : S_i \rightarrow S_i$ be the induced maps representing ϕ . Following the proof of Theorem 4.4, for $i \gg 0$, either $f_i : S_i \rightarrow S_i$ contains a noncontractible proper subgraph K_i that is f_i -invariant up to homotopy or all folds in $S_i \rightarrow S_j$ for $j > i$ are supported in some forest of S_i . In the former case, we get a reduction for ϕ and we are done. In the latter case, we can find a ϕ -periodic simplex $\sigma(S_j)$ in $SCV(F)$ for some $j > i$. Fix $j > i$ and $k \geq 1$ such that $\sigma(S_j) \cdot [\phi]^k = \sigma(S_j)$.
- (5) By Lemma 3.2, f_j^k is homotopic to an immersion $g : S_j \rightarrow S_j$. Collapse any maximal g -invariant forest if necessary and assume the immersion g has no invariant forest. If g is not irreducible, then we have a reduction for ϕ^k and so ϕ is not irreducible by Corollary 4.6. Assume the immersion g is irreducible.
- (6) By Corollary 2.8 again, either some Whitehead graph of g is disconnected so ϕ^k is reducible or g is clean. In the latter case, we get ϕ^k is irreducible by Theorem 4.5.

A careful analysis of Step (4) is required to understand the effectiveness of the algorithm.

5. SUBGROUPS INVARIANT UNDER IRREDUCIBLE ENDOMORPHISMS

In this section, we generalize a result by Bestvina-Feighn-Handel [2, Proposition 2.4] and I. Kapovich [12, Proposition 4.2] that characterizes the finitely generated subgroups of F that support a *leaf of the lamination* of an irreducible automorphism.

Definition 5.1. Given an expanding irreducible train track map $f : \Gamma \rightarrow \Gamma$, the **lamination of f** , denoted by $\Lambda(f)$, is defined by iterated neighbourhoods of f -periodic non-vertex points. Precisely, suppose $x \in \text{Int}(e)$ for some $e \in E(\Gamma)$ and $k \geq 1$ are such that $f^k(x) = x$. Then a **leaf of the lamination of f** , is the isometric immersion $\gamma_x : \mathbb{R} \rightarrow \Gamma$ such that $\gamma_x(0) = x$ and $f^k(\gamma_x(r)) = \gamma_x(\lambda_f^k \cdot r)$ for all $r \in \mathbb{R}$, unique up to orientation.

$$\Lambda(f) = \{\gamma_x : x \text{ is an } f\text{-periodic non-vertex point.}\}$$

For any integer $m \geq 1$, f^m -periodic points are f -periodic and vice-versa. So it follows that $\Lambda(f) = \Lambda(f^m)$ for any $m \geq 1$.

We now address the difficulty that arises when generalizing Bestvina-Feighn-Handel and I. Kapovich's results. Any automorphism $\phi : F \rightarrow F$ permutes the finite index subgroups with the same index; so given any finite index subgroup H' , there exists $i \geq 1$ such that $\phi^i(H') = H'$. This fact is used to lift a clean map representing ϕ to a clean map on the Stallings subgroup graph $S(H')$. However, this fails when dealing with nonsurjective injective endomorphisms, i.e., there is no reason why some ϕ -iterate of H' must be a subgroup of H' . The next lemma gives us a way of getting around this failure. The key observation is to look at backward iteration: when ϕ is injective, then pre-images of finite index subgroups are finite index subgroups with the same index or less.

Lemma 5.2. *Let $\phi : F \rightarrow F$ be an injective endomorphism of a free group F . If $H' \leq F$ is a finite index subgroup, then there exist $j > k \geq 0$ such that $\phi^{-k}(H') = \phi^{-j}(H')$.*

Furthermore, for $K = \phi^{-k}(H')$, there is an induced set bijection $\varphi : F/K \rightarrow F/K$ such that the following diagram commutes:

$$\begin{array}{ccc} F & \xrightarrow{\phi^{j-k}} & F \\ \downarrow \pi & & \downarrow \pi \\ F/K & \xrightarrow{\varphi} & F/K \end{array}$$

Proof. Set $H_k = \phi^{-k}(H')$ for $k \geq 0$. Then $\phi^k(H_k) = \phi^k(F) \cap H'$. Since H' has finite index in F , then $\phi^k(H_k) = \phi^k(F) \cap H'$ has finite index in $\phi^k(F)$. In fact, $[F : H'] \geq [\phi^k(F) : \phi^k(H_k)]$. But $\phi^k : F \rightarrow \phi^k(F)$ is an isomorphism that maps $H_k \leq F$ to $\phi^k(H_k) \leq \phi^k(F)$. Thus, $[\phi^k(F) : \phi^k(H_k)] = [F : H_k]$ and H_k has finite index in F bounded by $[F : H']$ for all $k \geq 0$. As there are only finitely many subgroups with index bounded by $[F : H']$, there must exist $j > k \geq 0$ such that $H_k = H_j$, i.e., $\phi^{-k}(H') = \phi^{-j}(H')$.

Let $K = \phi^{-k}(H')$ and $\pi : F \rightarrow F/K$ be the (left) coset projection map. The function $\pi \circ \phi^{j-k}$ factors through π if and only if $\phi^{j-k}(K) \leq K$, or equivalently, $K \leq \phi^{k-j}(K)$, and the induced function $\varphi : F/K \rightarrow F/K$ is an injection if and only if $\phi^{k-j}(K) \leq K$. By construction, $\phi^{k-j}(K) = K$, i.e., both conditions are satisfied, so $\pi \circ \phi^{j-k} = \varphi \circ \pi$ where φ is a bijection since it is an injection of the finite set F/K into itself. \square

The main result of the section follows:

Proposition 5.3. *Suppose $\phi : F \rightarrow F$ is injective and represented by a clean map. If $H \leq F$ is a finitely generated subgroup such that $\phi(H) \leq H$ and H contains a ϕ -expanding conjugacy class, then*

$$[\phi^k(F) : \phi^k(F) \cap H] < \infty \quad \text{for some } k \geq 0.$$

Proof. Suppose $\phi(H) \leq H$ for some finitely generated group $H \leq F$. Let $f : \Gamma \rightarrow \Gamma$ be the given clean map and ρ be the f -expanding immersed loop in Γ that lifts to a loop in $S(H)$. The invariance $\phi(H) \leq H$ implies the loops $[f^k(\rho)]$ lift to loops in $S(H)$ for all $k \geq 1$. As ρ is f -expanding and f is a clean map, length of $[f^k(\rho)]$ grows arbitrarily with k while the number of f -illegal turns in $[f^k(\rho)]$ remains bounded. Thus, for some $k \geq 1$, the loop $[f^k(\rho)]$ will contain an f -legal subpath longer than the critical constant. Therefore, there is a nontrivial subpath s of ρ such that $f^k(s)$ is a subpath of $[f^k(\rho)]$ for all $k \geq 1$. As f is clean, some f -iterate of s maps onto Γ .

Let x be an f -periodic non-vertex point of ρ and $\gamma_x : \mathbb{R} \rightarrow \Gamma$ be the corresponding leaf of $\Lambda(f)$. For all real $r > 0$, the path $\gamma_x([-r, r])$ is a subpath of $[f^k(\rho)]$ for some $k \geq 1$ since x is contained in some f -iterate of s . Thus $\gamma_x|_{[-r, r]}$ has a lift $\gamma_{x,r} : [-r, r] \rightarrow S(H)$. There are only finitely many preimages of x in $S(H)$, so after passing to an unbounded increasing subsequence r_m , we can assume the sequence $(\gamma_{x,r_m}(0))_{m=1}^\infty$ is constant. By uniqueness of lifts, $\gamma_{x,r_{m+1}}$ is an extension of γ_{x,r_m} for $m \geq 1$. The limit immersion $\gamma_{x,\infty} : \mathbb{R} \rightarrow S(H)$ is a lift of γ_x and we say $S(H)$ supports a leaf of $\Lambda(f)$.

As H is finitely generated, $S(H)$ is a finite graph and we can add edges to $S(H)$ if necessary to extend the immersion $S(H) \rightarrow \Gamma$ to a finite cover $S(H') \rightarrow \Gamma$ corresponding to $H \leq H' \leq F$. Thus $[F : H'] < \infty$ and we can apply Lemma 5.2 to get $j > k \geq 0$ such that $\phi^{-k}(H') = \phi^{-j}(H')$. Set $K = \phi^{-k}(H')$. Then $\phi^k(K) = \phi^k(F) \cap H'$ has finite index in $\phi^k(F)$. Recalling the diagram preceding Lemma 4.3, the graph $S(\phi^k(F)) = S_k$ and the cover corresponding to $\phi^k(K) \leq \phi^k(F)$ lie in the following commutative diagram:

$$\begin{array}{ccc}
 S(\phi^k(K)) & & S(\phi^k(K)) \\
 \downarrow p & & \downarrow p \\
 S(\phi^k(F)) & \xrightarrow{g} & S(\phi^k(F)) \\
 \downarrow v_k & \nearrow h_k & \downarrow v_k \\
 \Gamma & \xrightarrow{f^k} & \Gamma
 \end{array}$$

where $g : S(\phi^k(F)) \rightarrow S(\phi^k(F))$ is clean by Lemma 4.3 and $p : S(\phi^k(K)) \rightarrow S(\phi^k(F))$ is a finite cover. The map h_k maps f^k -periodic points to g -periodic points and v_k maps g -periodic points to f^k -periodic points. This allows us to identify $\Lambda(f)$ with $\Lambda(g)$.

Let $\Delta_H = S(\phi^k(F) \cap H)$. This is a noncontractible graph since $\phi(H) \leq H$. Note that $S(\phi^k(K))$ is the pullback of $S(\phi^k(F)) \rightarrow \Gamma$ and $S(H') \rightarrow \Gamma$ and Δ_H is the pullback of $S(\phi^k(F)) \rightarrow \Gamma$ and $S(H) \rightarrow \Gamma$. As $S(H)$ is a subgraph of $S(H')$, we have that Δ_H is a subgraph of $S(\phi^k(K))$. The graphs fit in this commutative diagram:

$$\begin{array}{ccccc}
 \Delta_H & \hookrightarrow & S(\phi^k(K)) & \xrightarrow{p} & S(\phi^k(F)) \\
 \downarrow & & \downarrow & & \downarrow v_k \\
 S(H) & \hookrightarrow & S(H') & \longrightarrow & \Gamma
 \end{array}$$

Since $S(\phi^k(F))$ and $S(H)$ support a leaf of $\Lambda(f)$, it follows that their pullback Δ_H supports a leaf of $\Lambda(f)$. Following the identification $\Lambda(f) \cong \Lambda(g)$, we have that Δ_H supports a leaf of $\Lambda(g)$. Lemma 5.2 says ϕ^{j-k} induces a permutation of $F/K \cong \phi^k(F)/\phi^k(K)$. Therefore, the map $g^{j-k} : S(\phi^k(F)) \rightarrow S(\phi^k(F))$ lifts to a map $\hat{g} : S(\phi^k(K)) \rightarrow S(\phi^k(K))$ with the property: if $g^{j-k}(x) = x$, then \hat{g} permutes the elements of $p^{-1}(x)$.

The rest of the argument follows that of Bestvina-Feighn-Handel [2, Lemma 2.1]. We include the details for completeness; see also [12, Proposition 4.2].

Claim. *The map $\hat{g} : S(\phi^k(K)) \rightarrow S(\phi^k(K))$ is clean.*

Proof. Let $\{a', b'\}$ be a turn at a vertex $v \in S(\phi^k(K))$ such that its projection under p , $\{a, b\}$, is an edge of the Whitehead graph of g^{j-k} at $p(v)$. Since g^{j-k} is a clean map, we can replace it with an iterate and assume $g^{j-k}(a) = \dots ab \dots$. So a contains a g^{j-k} -fixed point x and, consequently, \hat{g} permutes the lifts $p^{-1}(x)$. Replace g^{j-k} with an iterate if necessary and assume \hat{g} fixes $p^{-1}(x)$ and let $x' \in p^{-1}(x)$ be the lift of x in a' . Then x' is a \hat{g} -fixed point and $\hat{g}(a') = \dots a'b' \dots$. An identical argument shows that $\hat{g}(b') = \dots a'b' \dots$.

after passing to a power if necessary. Thus the Whitehead graph at v with respect to \hat{g} is isomorphic to the Whitehead graph at $p(v)$ with respect to g^{j-k} . So the Whitehead graphs of $S(\phi^k(K))$ with respect to \hat{g} are connected and \hat{g} is a train track as the turns in $\hat{g}(e')$ are lifts of the turns in $g^{j-k}(p(e'))$ for any edge e in $S(\phi^k(K))$. It remains to show that g is irreducible.

Since the Whitehead graphs of $S(\phi^k(K))$ are connected, for any edges a, b in $S(\phi^k(K))$, there is a sequences of turns $\{\epsilon_1, \epsilon_2\}, \{\epsilon_3, \epsilon_4\}, \dots, \{\epsilon_{2l-1}, \epsilon_{2l}\}$ such that each turn is an edge of the corresponding Whitehead graph, $\epsilon_{2m}, \epsilon_{2m+1}$ are half-edges of the same edge for $1 \leq m < l$, and $\epsilon_1, \epsilon_{2l}$ are half-edges of a, b respectively. Let $a = e_0, e_1, \dots, e_{l-1}, e_l = b$ the corresponding sequence of edges. By the previous paragraph, some iterate of \hat{g} maps e_m to e_{m+1} for $0 \leq m < l$. By induction, some iterate of \hat{g} maps a to b . As a and b were arbitrary, \hat{g} is an irreducible train track. \square

Claim. *The subgraph $\Delta_H \subset S(\phi^k(K))$ is not proper, i.e., $\Delta_H = S(\phi^k(K))$.*

Proof. Recall that a leaf of $\Lambda(g) = \Lambda(g^{j-k})$ is constructed by iterating neighbourhoods of some g^{j-k} -periodic non-vertex point. Suppose $x \in \text{Int}(e)$ for some edge e in $S(\phi^k(F))$ and $l \geq 1$ are such that $g^{(j-k)l}(x) = x$ and $g^{(j-k)l}$ lifts to a map $\hat{g}^l : S(\phi^k(K)) \rightarrow S(\phi^k(K))$ that fixes $p^{-1}(x)$ pointwise. Then the lift of $\gamma_x \in \Lambda(g^{j-k})$ to Δ_H can be constructed by using \hat{g}^l to iterate a neighbourhood of some $x' \in p^{-1}(x) \cap \Delta_H$. Let e' be the edge containing x' . Since \hat{g} is clean, $\hat{g}^m(e')$ surjects onto the whole graph $S(\phi^k(K))$ for some $m \geq 1$ and any iterated neighbourhood of x' shall eventually cover $S(\phi^k(K))$. Therefore, $S(\phi^k(K)) \subset \Delta_H$. \square

So the natural map $\Delta_H \rightarrow S(\phi^k(F))$ is a finite cover and $[\phi^k(F) : \phi^k(F) \cap H] < \infty$. \square

We shall now use it to give sufficient conditions for an endomorphism to be fully irreducible. The reader only interested in our hyperbolicity result can skip to the next section.

Proposition 5.4. *If $\phi : F \rightarrow F$ is injective, primitively atoroidal, represented by a clean map, and its image $\phi(F)$ is not contained in a proper free factor, then ϕ is fully irreducible.*

Proof. Let $A \leq F$ be a ϕ^n -invariant free factor for some $n \geq 1$ and assume A has minimal rank (for all n). Then $i_g \circ \phi^n(A) \leq A$ for some inner automorphism i_g . If A is cyclic then it is generated by an element in a ϕ -expanding conjugacy class – the definition of being *primitively atoroidal*. Otherwise, the minimality assumption implies $i_g \circ \phi^n|_A$ is fully irreducible. In both cases, A contains a ϕ -expanding conjugacy class. As ϕ is represented by a clean map, so is $i_g \circ \phi^n$. By Proposition 5.3, there exists $k \geq 0$ such that $(i_g \circ \phi^n)^k(F) \cap A$ has finite index in $(i_g \circ \phi^n)^k(F)$. It remains to show $A = F$, hence, ϕ is fully irreducible.

Since A is a free factor of F , the intersection $(i_g \circ \phi^n)^k(F) \cap A$ is a free factor of $(i_g \circ \phi^n)^k(F)$. The only finite index free factor of a finitely generated free group is the free group itself. Thus $(i_g \circ \phi^n)^k(F) \cap A = (i_g \circ \phi^n)^k(F)$ and $(i_g \circ \phi^n)^k(F) \leq A$.

Recall that $\phi(F)$ is not contained in a proper free factor. Suppose $\phi^{l-1}(F)$ is not contained in a proper free factor of F for some $l \geq 2$. Let $K \leq F$ be a free factor such that $\phi^l(F) \leq K$. Then $\phi(\phi^{l-1}(F)) = \phi^l(F) \leq K \cap \phi^{l-1}(F)$ and the intersection is a free factor

of $\phi^{l-1}(F)$. As ϕ^{l-1} is injective, we have $\phi|_{\phi^{l-1}(F)}$ is conjugate to ϕ . Therefore, $\phi^l(F)$ is not contained in a proper free factor of $\phi^{l-1}(F)$ and $K \cap \phi^{l-1}(F) = \phi^{l-1}(F)$. Therefore $\phi^{l-1}(F) \leq K$. The induction hypothesis is that $\phi^{l-1}(F)$ is not contained in a proper free factor of F , hence $K = F$. So $\phi^l(F)$ is not contained in a proper free factor of F .

By induction, $\phi^n(F)$ is not contained in a proper free factor of F . We also have $i_g \circ \phi^n(F)$ is not contained in a proper free factor of F since i_g is an automorphism. By the same induction argument, $(i_g \circ \phi^n)^k(F)$ is not contained in a proper free factor of F . Therefore, $(i_g \circ \phi^n)^k(F) \leq A$ implies $A = F$ and ϕ is fully irreducible. \square

Corollary 5.5. *Let $\phi : F \rightarrow F$ be an injective endomorphism. Then ϕ is fully irreducible if and only if $\phi : F \rightarrow F$ is primitively atoroidal, represented by a clean map, and its image $\phi(F)$ is not contained in a proper free factor.*

This corollary extends the characterisation of fully irreducible automorphisms due to I. Kapovich [13, Theorem 1.2], which in turn was motivated by Catherine Pfaff's criterion for irreducibility [17, Theorem 4.1]. Comparing with Corollary 4.6, it seems that the condition on the image $\phi(F)$ is redundant.

6. IRREDUCIBLE NONSURJECTIVE ENDOMORPHISMS ARE HYPERBOLIC

The goal of this final section to prove that the mapping tori of irreducible nonsurjective endomorphisms are word-hyperbolic.

Definition 6.1. Let $\phi : F \rightarrow F$ be an injective endomorphism. Then the **ascending HNN extension/mapping torus** of ϕ is given by the presentation:

$$F*_\phi = \langle F, t \mid t^{-1}xt = \phi(x), \forall x \in F \rangle$$

Thurston's hyperbolization theorem gives the correspondence between the geometry of 3-manifolds that fiber over a circle and the dynamics of their monodromies [21] and Brinkmann generalized this to free-by-cyclic groups $F \rtimes \mathbb{Z}$ [7]. The following theorem, the main result of [16], is a partial generalization to ascending HNN extensions $F*_\phi$.

Theorem 6.2 ([16, Theorem 6.3]). *Suppose $\phi : F \rightarrow F$ is represented by an immersion. Then $F*__\phi$ is word-hyperbolic if and only if there are no $d, n \geq 1$ and $1 \neq a \in F$ such that $\phi^n(a)$ is conjugate to a^d in F .*

Remarkably, this theorem combined with Theorem 4.4 and Proposition 5.3 implies that irreducible nonsurjective endomorphisms have word-hyperbolic mapping tori.

Theorem 6.3. *If $\phi : F \rightarrow F$ is nonsurjective and irreducible, then $F*__\phi$ is word-hyperbolic.*

Proof. By Theorem 4.4, we can represent ϕ by a clean immersion $f : \Gamma \rightarrow \Gamma$ and every nontrivial conjugacy class is ϕ -expanding. Suppose $F*__\phi$ were not word-hyperbolic. By Theorem 6.2, there exists a nontrivial element $a \in F$, an element $g \in F$, and integers $d, n \geq 1$ such that $\phi^n(a) = ga^d g^{-1}$. If we let $H = \langle a \rangle$, then $i_g \circ \phi^n(H) \leq H$. By Proposition 5.3, there is a $k \geq 0$ such that $(i_g \circ \phi^n)^k(F) \cap H$ has finite index in $(i_g \circ \phi^n)^k(F)$. But this is a contradiction as H cannot be cyclic and have finite index intersection in a noncyclic free group. Therefore, $F*__\phi$ must be word-hyperbolic. \square

On the other hand, there are fully irreducible automorphisms whose corresponding free-by-cyclic groups are not word-hyperbolic. In this case, Bestvina-Handel showed that the automorphisms are induced by pseudo-Anasov homeomorphisms on once-punctured surface [4, Proposition 4.5]. By Thurston's hyperbolization theorem, the free-by-cyclic groups are fundamental groups of hyperbolic 3-manifolds that fiber over a circle.

REFERENCES

- [1] R. B. Bapat and T. E. S. Raghavan. *Nonnegative matrices and applications*, volume 64 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1997.
- [2] Mladen Bestvina, Mark Feighn, and Michael Handel. Laminations, trees, and irreducible automorphisms of free groups. *Geom. Funct. Anal.*, 7(2):215–244, 1997.
- [3] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for $\text{Out}(F_n)$. I. Dynamics of exponentially-growing automorphisms. *Ann. of Math. (2)*, 151(2):517–623, 2000.
- [4] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [5] Robert Bieri, Walter D. Neumann, and Ralph Strebel. A geometric invariant of discrete groups. *Invent. Math.*, 90(3):451–477, 1987.
- [6] Robert Bieri, Walter D. Neumann, and Ralph Strebel. A geometric invariant of discrete groups. *Invent. Math.*, 90(3):451–477, 1987.
- [7] Peter Brinkmann. Hyperbolic automorphisms of free groups. *Geom. Funct. Anal.*, 10(5):1071–1089, 2000.
- [8] Marc Culler and Karen Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.*, 84(1):91–119, 1986.
- [9] Warren Dicks and Enric Ventura. *The group fixed by a family of injective endomorphisms of a free group*, volume 195 of *Contemporary Mathematics*. American Mathematical Society, Providence, RI, 1996.
- [10] Spencer Dowdall, Ilya Kapovich, and Christopher J. Leininger. Dynamics on free-by-cyclic groups. *Geom. Topol.*, 19(5):2801–2899, 2015.
- [11] Spencer Dowdall, Ilya Kapovich, and Christopher J. Leininger. Endomorphisms, train track maps, and fully irreducible monodromies. *Groups Geom. Dyn.*, 11(4):1179–1200, 2017.
- [12] Ilya Kapovich. Algorithmic detectability of iwip automorphisms. *Bull. Lond. Math. Soc.*, 46(2):279–290, 2014.
- [13] Ilya Kapovich. Detecting fully irreducible automorphisms: a polynomial time algorithm. *Exp. Math.*, 28(1):24–38, 2019. With an appendix by Mark C. Bell.
- [14] Gilbert Levitt and Martin Lustig. Irreducible automorphisms of F_n have north-south dynamics on compactified outer space. *J. Inst. Math. Jussieu*, 2(1):59–72, 2003.
- [15] Jean Pierre Mutanguha. Irreducibility of a free group endomorphism is a mapping torus invariant. *in preparation*.
- [16] Jean Pierre Mutanguha. Hyperbolic Immersions of Free Groups. *Groups Geom. Dyn.*, to appear. arxiv:1809.04761.
- [17] Catherine Pfaff. Ideal Whitehead graphs in $\text{Out}(F_r)$ II: the complete graph in each rank. *J. Homotopy Relat. Struct.*, 10(2):275–301, 2015.
- [18] Patrick Reynolds. Dynamics of Irreducible Endomorphisms of F_n . *ArXiv e-prints*, August 2010. arxiv:1008.3659.
- [19] John R. Stallings. On fibering certain 3-manifolds. In *Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)*, pages 95–100. Prentice-Hall, Englewood Cliffs, N.J., 1962.
- [20] John R. Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [21] William P. Thurston. Three-dimensional manifolds, Kleinian groups and hyperbolic geometry. *Bull. Amer. Math. Soc. (N.S.)*, 6(3):357–381, 1982.

- [22] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):417–431, 1988.
- [23] Karen Vogtmann. Automorphisms of free groups and outer space. In *Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part I (Haifa, 2000)*, volume 94, pages 1–31, 2002.
- [24] Karen Vogtmann. Contractibility of outer space: reprise. In *Hyperbolic geometry and geometric group theory*, volume 73 of *Adv. Stud. Pure Math.*, pages 265–280. Math. Soc. Japan, Tokyo, 2017.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ARKANSAS, FAYETTEVILLE, AR

Web address: <https://mutanguha.com>

Email address: jpmutang@uark.edu